

DO NOT STOP taking medications prescribed by your doctor in order to donate blood. Donating while taking these drugs could have a negative effect on your health or on the health of the recipient of your blood. PLEASE TELL US IF YOU:

TBC Medication Deferral List							
Are being treated with... Testosterone Agents	also called ... Testosterone, Andro, Depandro 100, Depo-Testosterone, Cypionate, Testosterone Enanthate			anytime in the last ... 2 weeks			
AABB Medication Deferral List – v4.0							
Are being treated with any of the following types of medications....	or have taken...		which is also called...	anytime in the last....			
Deferral for Platelet Donors Only Anti-platelet agents (usually taken to prevent stroke or heart attack)	Feldene		piroxicam	2 days			
	Effient		prasugrel	3 days			
	Brilinta		ticagrelor	7 days			
	Plavix		clopidogrel	14 days			
	Ticlid		ticlopidine				
	Zontivity		vorapaxar	1 month			
Anticoagulants or “blood thinners” (usually taken to prevent blood clots in the legs and lungs and to prevent strokes)	Arixtra		fondaparinux	2 days			
	Eliquis		apixaban				
	Fragmin		dalteparin				
	Lovenox		enoxaparin				
	Pradaxa		dabigatran				
	Savaysa		edoxaban				
	Xarelto		rivaroxaban				
	Coumadin, Warfilone, Jantoven		warfarin	7 days			
	Heparin, low-molecular-weight heparin						
Acne treatment	Accutane, Amnesteem, Absorica, Claravis, Myorisan, Sotret, Zenatane		isotretinoin	1 Month			
Multiple myeloma	Thalomid Revlimid		thalidomide lenalidomide				
Rheumatoid arthritis	Rinvoq		upadacitinib				
Hair loss remedy	Propecia		finasteride				
Prostate symptoms	Proscar		finasteride				
	Avodart, Jalyn		dutasteride	6 Months			
Immunosuppressant	Cellcept		mycophenolate mofetil	6 weeks			
HIV prevention (also known as PrEP or PEP)	Any medication taken by mouth (oral) to prevent HIV.	Truvada	emtricitabine and tenofovir disoproxil fumarate	4 months			
		Descovy	emtricitabine and tenofovir alafenamide				
	Injectable HIV prevention	Apretude	cabotegravir	24 months			
Basal cell skin cancer	Erivedge, Odomzo		vismodegib, sonidegib	24 months			
Relapsing multiple sclerosis	Aubagio		teriflunomide				
Rheumatoid arthritis	Arava		leflunomide				
Hepatitis exposure	Hepatitis B Immune Globulin		HBIG	12 months			
Experimental Medication or Unlicensed (Experimental) Vaccine							
Psoriasis	Soriatane		acitretin	36 months			
	Tegison		etretinate	Ever			
HIV treatment - Any medication to treat HIV. May also be called antiretroviral therapy (ART)							
Growth hormone from human pituitary glands (no longer available in US)							

Material educativo para donantes de sangre

¡DEBE LEER ESTO ANTES DE DONAR!

- **Sus respuestas precisas y honestas son fundamentales para la seguridad de los pacientes que reciben transfusiones de sangre.**
- Cada pregunta es necesaria para evaluar completamente la seguridad de su donación.
- Según lo exige la normativa, le indicamos que no done sangre si tiene un factor de riesgo.
- Si no entiende una pregunta, pida ayuda al personal del centro de donación de sangre.
- SUS RESPUESTAS SON CONFIDENCIALES.

Para determinar si usted cumple con los requisitos para donar, haremos lo siguiente:

- Le preguntaremos sobre su salud y los medicamentos que está tomando o ha tomado.
- Le preguntaremos si ha viajado o vivido en otros países.
- Le preguntaremos sobre su riesgo de contraer infecciones que pueden transmitirse por la sangre, especialmente el VIH (que es el virus que causa el SIDA) y la hepatitis viral.
- Tomaremos su presión arterial, temperatura y pulso.
- Le tomaremos una muestra de sangre para verificar que su recuento sanguíneo sea aceptable antes de donar.

Si usted cumple con los requisitos para donar, haremos lo siguiente:

- Le limpiaremos el brazo con un antiséptico (díganos si tiene alguna alergia en la piel).
- Usaremos una aguja y un juego de tubos estériles para recolectar su sangre.

NUNCA reutilizamos una aguja o un juego de tubos.

QUÉ SUCEDE DESPUÉS DE SU DONACIÓN

Para proteger a los pacientes, su sangre se analiza para detectar hepatitis B y C, VIH, sífilis y otras infecciones. Si sus análisis de sangre dan positivo, no se le dará a ningún paciente. Se le notificará sobre cualquier resultado positivo de la prueba que pueda afectar su elegibilidad para donar en el futuro. Hay ocasiones en las que no se analiza su sangre. Si esto ocurre, es posible que no reciba ninguna notificación. El centro de donación de sangre no divulgará los resultados de su prueba sin su permiso por escrito, a menos que sea exigido por la ley (p. ej., al Departamento de Salud).

ELEGIBILIDAD DEL DONANTE – INFORMACIÓN ESPECÍFICA

Ciertas enfermedades infecciosas, como el VIH y la hepatitis, se pueden propagar a través de:

- El contacto sexual
- Otras actividades que aumenten el riesgo
- Transfusión de sangre

Le haremos preguntas específicas sobre el contacto sexual y otras actividades que pueden aumentar el riesgo de estas infecciones.

Qué entendemos por "contacto sexual"

Las palabras “tener contacto sexual con” y “sexo” se utilizan en algunas de las preguntas que le haremos. Estas preguntas se aplican a todas las actividades a continuación, ya sea que se usaran o no medicamentos, condones u otra protección para prevenir infecciones o embarazos:

- Sexo vaginal (contacto entre el pene y la vagina)
- Sexo oral (boca o lengua en la vagina, el pene o el ano de alguien)
- Sexo anal (contacto entre el pene y el ano)

Una “nueva pareja sexual” incluye los siguientes ejemplos:

- Tener sexo con alguien por primera vez
O
- Haber tenido relaciones sexuales con alguien en una relación que terminó en el pasado y haber vuelto a tener relaciones sexuales con esa persona en los últimos 4 meses.

Factores de riesgo del VIH y la hepatitis

El VIH y la hepatitis se transmiten principalmente por contacto sexual con una persona infectada O por compartir agujas o jeringas que una persona infectada usa para inyectarse drogas.

NO DONE si usted:

- **ALGUNA VEZ ha tomado algún medicamento para tratar la infección por VIH.**
- **Está tomando algún medicamento para prevenir la infección por VIH. Estos medicamentos se pueden llamar PrEP, PEP, TRUVADA, DESCOVY, APRETUDE o muchos otros nombres.**

Los medicamentos antirretrovirales aprobados por la FDA son seguros y eficaces para prevenir la transmisión sexual del VIH. Sin embargo, estos medicamentos antirretrovirales no eliminan por completo el virus del cuerpo, y la sangre donada aún puede potencialmente transmitir la infección por el VIH al receptor de la transfusión.

NO DEJE DE TOMAR NINGÚN MEDICAMENTO RECETADO PARA DONAR SANGRE, INCLUSO LOS MEDICAMENTOS DE PrEP y PEP.

NO DONE si usted:

- **ALGUNA VEZ** ha tenido una prueba positiva para la infección por VIH.
- **En los últimos 4 meses:**
 - Ha tenido contacto sexual con una nueva pareja **y** ha tenido sexo anal.
 - Ha tenido contacto sexual con más de una pareja **y** ha tenido sexo anal.
 - Ha tenido contacto sexual con cualquier persona que alguna vez haya tenido una prueba positiva de infección por VIH.
 - Ha recibido dinero, drogas u otro pago por sexo.
 - Usó agujas para inyectarse drogas, esteroides o cualquier cosa no recetada por su médico.
 - Ha tenido contacto sexual con cualquier persona que haya recibido dinero, drogas u otro pago por sexo, **o** que haya usado aguj

Blood Donor Educational Material

YOU MUST READ THIS BEFORE YOU DONATE!

- Your **accurate and honest** responses are critical to the safety of patients who receive blood transfusions.
- Each question is necessary to fully evaluate the safety of your donation.
- As required by regulations, we are instructing you not to donate blood if you have a risk factor.
- If you don't understand a question, ask the blood center staff for assistance.
- YOUR RESPONSES ARE CONFIDENTIAL.

To determine if you are eligible to donate, we will:

- Ask about your health and medications you are taking or have taken.
- Ask if you have traveled to or lived in other countries.
- Ask about your risk for infections that can be transmitted by blood – especially HIV (which is the virus that causes AIDS), and viral hepatitis.
- Take your blood pressure, temperature, and pulse.
- Take a blood sample to be sure your blood count is acceptable before you donate.

If you are eligible to donate, we will:

- Clean your arm with an antiseptic (Tell us if you have any skin allergies).
- Use a sterile needle and tubing set to collect your blood.

We NEVER reuse a needle or tubing set.

WHAT HAPPENS AFTER YOUR DONATION

To protect patients, your blood is tested for hepatitis B and C, HIV, syphilis, and other infections. If your blood tests positive, it will not be given to a patient. You will be notified about any positive test result which may affect when you are eligible to donate in the future. There are times when your blood is not tested. If this occurs, you may not receive any notification. The blood center will not release your test results without your written permission unless required by law (e.g., to the Health Department).

DONOR ELIGIBILITY – SPECIFIC INFORMATION

Certain infectious diseases, such as HIV and hepatitis, can be spread through:

- Sexual contact
- Other activities that increase risk
- Blood transfusion

We will ask specific questions about sexual contact and other activities that may increase risk for these infections.

What do we mean by “sexual contact?”

The words “have sexual contact with” and “sex” are used in some of the questions we will ask you. These questions apply to all of the activities below, whether or not medications, condoms or other protection were used to prevent infection or pregnancy:

- Vaginal sex (contact between penis and vagina)
- Oral sex (mouth or tongue on someone's vagina, penis, or anus)
- Anal sex (contact between penis and anus)

A “new sexual partner” includes the following examples:

- Having sex with someone for the first time
OR
- Having had sex with someone in a relationship that ended in the past, and having sex again with that person in the last 4 months.

HIV/Hepatitis risk factors

HIV and hepatitis are spread mainly by sexual contact with an infected person OR by sharing needles or syringes used by an infected person to inject drugs.

DO NOT DONATE if you:

- Have **EVER** taken any medication **to treat HIV** infection.
- Are taking any medication **to prevent HIV** infection. These medications may be called: **PrEP, PEP, TRUVADA, DESCOVY, APRETUDE** or many other names.

FDA-approved antiretroviral drugs are safe and effective in preventing sexual transmission of HIV. However, these antiretroviral drugs do not fully eliminate the virus from the body, and donated blood can potentially still transmit HIV infection to a transfusion recipient.

DO NOT STOP TAKING ANY PRESCRIBED MEDICATIONS IN ORDER TO DONATE BLOOD, INCLUDING PrEP and PEP MEDICATIONS.

DO NOT DONATE if you:

- Have **EVER** had a positive test for HIV infection.
- **In the past 4 months:**
 - Have had sexual contact with a new partner **and** have had anal sex.
 - Have had sexual contact with more than one partner **and** have had anal sex.
 - Have had sexual contact with anyone who has ever had a positive test for HIV infection.
 - Have received money, drugs, or other payment for sex.
 - Have used needles to inject drugs, steroids, or anything not prescribed by your doctor.
 - Have had sexual contact with anyone who has received money, drugs, or other payment for sex, **or** used needles to inject drugs, steroids, or anything not prescribed by their doctor.
 - Have had syphilis or gonorrhea or been treated for syphilis or gonorrhea.
- **In the past 12 months:**
 - Have been in juvenile detention, lockup, jail or prison for 72 hours or more consecutively.
- Have **EVER** had Ebola virus infection or disease.

DO NOT DONATE if you have these symptoms which can be present before you test positive for HIV:

- Fever
- Enlarged lymph glands
- Sore throat
- Rash

Your blood can transmit infections, including HIV, even if you feel well and all your tests are normal. Even the best tests cannot detect the virus for a period of time after you are infected.

DO NOT DONATE:

- If you think you may be at risk